



Prairie and Peregrine Falcon Occupancy and Productivity Monitoring at Pinnacles National Park

2016 Annual Report

Natural Resource Report NPS/SFAN/NRR—2017/1379



ON THE COVER

Prairie falcon fledgling, Discovery Wall, Pinnacles National Park, California.
NPS /GAVIN EMMONS

Prairie and Peregrine Falcon Occupancy and Productivity Monitoring at Pinnacles National Park

2016 Annual Report

Natural Resource Report NPS/SFAN/NRR—2017/1379

Gavin Emmons

National Park Service
Pinnacles National Park
5000 Highway 146
Paicines, California 95043

January 2017

U.S. Department of the Interior
National Park Service
Natural Resource Stewardship and Science
Fort Collins, Colorado

The National Park Service, Natural Resource Stewardship and Science office in Fort Collins, Colorado, publishes a range of reports that address natural resource topics. These reports are of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Report Series is used to disseminate comprehensive information and analysis about natural resources and related topics concerning lands managed by the National Park Service. The series supports the advancement of science, informed decision-making, and the achievement of the National Park Service mission. The series also provides a forum for presenting more lengthy results that may not be accepted by publications with page limitations.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and designed and published in a professional manner. This report received informal peer review by subject-matter experts who were not directly involved in the collection, analysis, or reporting of the data. Data in this report were collected and analyzed using methods based on established, peer-reviewed protocols and were analyzed and interpreted within the guidelines of the protocols.

Views, statements, findings, conclusions, recommendations, and data in this report do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the U.S. Government.

This report is available in digital format from the [San Francisco Bay Area Network Inventory and Monitoring website](#) and the [Natural Resource Publications Management website](#). To receive this report in a format optimized for screen readers, please email irma@nps.gov.

Please cite this publication as:

Emmons, G. 2016. Prairie and peregrine falcon occupancy and productivity monitoring at Pinnacles National Park: 2016 annual report. Natural Resource Report NPS/SFAN/NRR—2017/1379. National Park Service, Fort Collins, Colorado.

Contents

	Page
Figures.....	iv
Tables.....	iv
Appendices.....	iv
Executive Summary	v
Acknowledgments.....	vi
Introduction.....	7
Study Area and Field Methods	9
Sample Design.....	9
Field Methods.....	11
Monitoring Schedule	12
Data Management.....	12
Climbing Advisories.....	12
Results.....	14
Prairie Falcons	14
Occupied Territories	14
Annual Productivity.....	15
Phenology	20
Other Notes.....	20
Discussion	21
Conclusions, Management Implications and Recommendations	21
Prairie Falcons: Climbing Advisories	21
Human / Falcon Interactions and Nest Failures	22
Education Opportunities.....	22
Management Recommendations	22
Literature Cited	24

Figures

	Page
Figure 1. Core and non-core areas at Pinnacles National Park.	10
Figure 2. Setting up advisory sign (NPS/GAVIN EMMONS, 2014).	13
Figure 3. Core vs. non-core Pinnacles prairie falcon fledgling productivity, 1984–2016.	17

Tables

	Page
Table 1. Timing of nesting behavior of prairie falcons at Pinnacles National Park.	12
Table 2. 1984–2016 Pinnacles prairie falcon nesting productivity – core and non-core areas combined.	15
Table 3. 2016 Pinnacles prairie falcon breeding summary.	16
Table 4. 1984–2016 Pinnacles prairie falcon nesting productivity – core areas only.	18
Table 5. 1984–2016 Pinnacles prairie falcon nesting productivity – non-core areas only.	19

Appendices

	Page
Appendix A. 2016 nest phenology and success for prairie and peregrine falcons.	27
Appendix B. 2016 results and discussion for peregrine falcon occupancy and productivity.	29
Appendix C. Documentation of changes in data collection methods.	34
Appendix D. Public interest highlights.	35

Executive Summary

Pinnacles National Park (“Pinnacles”) provides a diverse habitat for numerous cliff-nesting raptors, including prairie falcons (*Falco mexicanus*) and peregrine falcons (*F. peregrinus*), as well as a spectacular array of summits and cliff-wall routes for rock-climbers. This monitoring program was established to determine long-term trends in the number of occupied territories and productivity of nesting prairie and peregrine falcons. The monitoring program grew out of a need to reduce potential disturbance that climbers and off-trail hikers may have on cliff-nesting raptors. This report summarizes the results from the 2016 breeding season and represents the 31st year of monitoring at the park, consistent with the standardized methods and procedures detailed in the Prairie Falcon Monitoring Protocol for Pinnacles National Monument (Emmons et al. 2011).

To monitor falcons, field technicians surveyed all potential nest sites three times per breeding season, with visits spaced 21–28 days apart. Nests determined to be active were revisited to confirm rearing of nestlings and fledging of young. In 2016, monitoring was conducted from 1 January 2016 until 1 July 2016, with a total of over 150 possible and active nest sites monitored during 710 observation hours.

Thirteen territorial pairs of prairie and peregrine falcons were documented this year with 8 pairs actively nesting. Six nests successfully fledged 22 young; two nests failed.

Acknowledgments

This program would not be as successful as it is without the eyes and ears of helpful Pinnacles employees. Therefore, I would like to thank the National Park Service employees for their help, encouragement, and passion for the raptors and wildlife diversity at Pinnacles. The many local climbers involved with Friends of Pinnacles also deserve my thanks for their ongoing support of resource protection and breeding raptors at the park and particularly their efforts to publicize and honor advisories in effect. I would also like to extend my appreciation to the park visitors, for their reports and observations on raptor sightings and for their appreciation and value of the importance of monitoring, managing, and protecting the nesting sites and breeding productivity of raptors in the park.

I would like to thank J. Belli, A. Schmidt, M. Shelley, A. Punzalan, K. Vanesky, M. Baird, J. Negreann, M. Clark, A. List, Y. Pohl, J. Scullen, E. Taketa, S. Robinson, and A. Welch for contributing valuable observations on raptor territories and pair behavior in the park. I greatly appreciated efforts by E. Cox for her excellent monitoring efforts this season, and for the invaluable support that M. Gnekow provided as a raptor monitoring volunteer in 2016. I am also grateful for B. Johnson's and P. Johnson's support and efforts, in tandem with D. George as the manager of the Inventory and Monitoring (I&M) Program, to keep the Pinnacles raptor monitoring program funded annually and on a permanent basis. S. Wakamiya also provided valuable assistance in her role as the I&M Program Data Manager revising the design and enhancing the functionality of the raptor monitoring database. D. Adams, D. George, and P. Johnson also provided recommendations and reviewer suggestions for the 2016 annual report, contributing to a concise and efficient document consistent with I&M standards.

The following staff also shared their experience, excitement, and observations of raptors with me throughout the season, granting me a more complete picture of raptor breeding and diversity at the park, and assisted in the effective management of raptor advisory areas: P. Johnson, B. Johnson, D. Ryan, J. Jones, L. Regan, and J. Donham.

Introduction

Pinnacles National Park (“Pinnacles”) is a National Park Service (NPS) unit located in the Gabilan Mountains of central California, and was legislatively converted from a national monument to a national park in January 2013. Pinnacles provides a diverse habitat for cliff-nesting raptor species, including sensitive species such as prairie falcons (*Falco mexicanus*), peregrine falcons (*F. peregrinus*), and golden eagles (*Aquila chrysaetos*). The dramatic landscapes, extensive trails, arrays of summits, and cliff-wall routes at Pinnacles are also used intensively for recreation by rock-climbers and hikers. Because prairie falcons nest in the Pinnacles cliffs and in sufficient density to track trends in reproduction over time, this species is the central focus of the monitoring program (Emmons et al. 2011). However, both prairie and peregrine falcons have nested at Pinnacles historically, are sensitive to human disturbance, are obligate cliff-nesters, and have identical nest phenology patterns. Because they occupy the same nest habitat and are direct competitors to prairie falcons, peregrine falcon monitoring is also documented in this report. Other raptor species in the park either nest in forested habitats or do not nest in sufficient densities within the park to warrant a similar level of monitoring effort.

Many scientific studies have documented the negative impacts of human disturbance of raptor nest and roost sites, and the resulting nest failures and territorial abandonment associated with these disturbances. Nesting raptor species at Pinnacles sensitive to human disturbance include prairie falcons (Fyfe and Olendorff 1976; Ogden and Hornocker 1977; Harmata et al. 1978; Sitter 1983; Steenhof 1998), peregrine falcons (particularly in remote locations: see Hickey 1942, 1969; Bond 1946; Steenhof 1998), golden eagles (Newton 1979, 1990; Scott 1985; Steidl et al. 1993; Watson 1997; Steenhof et al. 1997; Kochert et al. 1999), sharp-shinned hawks (*Accipiter striatus*; Delannoy and Cruz 1988), and long-eared owls (*Asio otus*; Marks 1986; Marti and Marks 1989; Bloom 1994).

Studies of prairie and peregrine falcon nest occupancy and productivity have also shown the species to be especially sensitive to human disturbance from mining (Becker and Ball 1981; Bednarz 1984), recreation (Boyce 1982), agriculture (USDI 1979), habitat destruction and nest site limitation (Becker and Ball 1981; Steenhof et al. 1997), and proximity to major roadways (Platt 1974; Boyce 1982).

The main sources of human disturbance of nesting falcons at Pinnacles are visitors who are rock-climbing and hiking on- and off-trail in the park. Scientific studies have consistently suggested that these recreational activities can be balanced against raptor nesting by establishing closure or advisory areas that act as buffers between human activity and raptor nesting during the breeding season (Fyfe and Olendorff 1976; Olsen and Olsen 1978; Becker and Ball 1981; Suter and Jones 1981; Porter et al. 1987; Holthuijzen et al. 1990; Cade et al. 1996; White et al. 2002). Raptor monitoring program survey data collected at Pinnacles support the establishment of climbing/hiking advisories in core areas (high visitor-use areas) each breeding season as a way to protect cliff-nesting raptor species from human disturbance.

The Prairie Falcon Monitoring Protocol for Pinnacles National Monument (Emmons et al. 2011) provides standardized methods and procedures for prairie and peregrine falcon monitoring at

Pinnacles and further details the program specifics. The program established two long-term monitoring objectives to:

- Track changes in the total numbers of territorial falcon pairs in core areas and non-core areas.
- Track changes in average annual productivity (young of year hatched/pair, young of year at banding age/pair, young of year fledged/pair) in core areas and non-core areas.

Core areas are locations at Pinnacles suitable for prairie and peregrine falcon cliff-nesting where climbing impacts could occur, based on the presence of historic climbing routes accessible to visitors. Non-core areas refer to all other areas within Pinnacles suitable for cliff-nesting. The core vs. non-core sampling design is detailed further in the Methods section.

A secondary benefit of the monitoring program is that a substantial amount of information can also be gathered on other raptor species at Pinnacles, particularly sensitive California species that may be impacted by human presence and disturbance in riparian habitats such as: golden eagles, Cooper's hawks (*Ac. cooperii*), sharp-shinned hawks, white-tailed kites (*Elanus leucurus*), and long-eared owls. Breeding data collected on other raptor species during the 2016 season will be documented in the Breeding Raptor Distribution and Nesting Phenology at Pinnacles National Park, 2012–2016 Report (Emmons in development).

Study Area and Field Methods

Pinnacles is located in the Gabilan Mountains of the central Coast Range of California. The national park encompasses 10,694 hectares (26,425 acres) with elevation ranging from 244 to 1,007 meters (800 to 3,304 feet). The climate is Mediterranean with hot, dry summers and cool, damp winters. Temperatures range from a mean of 5.2°C in December to 25.2°C in August (41.4° to 77.4°F). The average yearly rainfall is 42.3 cm (16.6 inches), with the majority of rainfall occurring from November to April (WRCC 2013).

Pinnacles provides a diverse range of habitat types for birds and other species. These habitats include volcanic rock formations and outcroppings, California mixed chaparral, pine-oak woodlands, grasslands, and riparian creek corridors.

Sample Design

The prairie and peregrine falcon monitoring is focused on core areas and non-core areas. Each core and non-core area is a potential nesting territory, and in this context the terms “area” and “territory” can be used interchangeably. Within a given year, prairie or peregrine falcons may actively defend one or more of these areas or territories. Core areas (Figure 1) are locations in Pinnacles that can support falcon cliff nesting, and where impacts to raptors due to rock climbing activities can occur based on historic rock-climbing use and access. Core area monitoring surveys are conducted through a census, because the area is sufficiently small to allow for complete coverage.

Non-core areas refer to all other areas within the park that can support prairie and peregrine falcon cliff nesting. The Prairie Falcon Monitoring Protocol for Pinnacles National Monument (Emmons et al. 2011) calls for sampling non-core areas on a rotating basis. For 2003–2016, non-core area monitoring has been conducted through a census along with core area censuses. This park-wide censusing of core and non-core areas has been possible because of comprehensive historical data on prairie and peregrine falcon nest sites gathered over the past 31 years, extensive monitoring experience of the raptor biologist, and supplemental raptor monitoring efforts by interns, volunteers, and other Pinnacles employees. In addition, GIS modeling completed in 2008 confirmed that all potential prairie and peregrine falcon nesting areas in the park have been surveyed annually during the past 14 years.




Figure 1. Core and non-core areas at Pinnacles National Park.

Field Methods

Survey methods followed the standard operating procedures detailed in the Prairie Falcon Monitoring Protocol for Pinnacles National Monument (Emmons et al. 2011).

Potential and established prairie and peregrine falcon territories in core and non-core areas were surveyed using Swarovski HD STS-80 and ATS-65 20–60x spotting scopes and Zeiss Victory FL 10×32 binoculars. Observations were made from the locations that provided the best view of an eyrie or a territory. Magellan Triton 500 and DeLorme Earthmate PN-60 GPS units were used to plot every observation point. Field data were documented with standardized datasheets and field notebooks and the data were entered into a Microsoft Access database (Appendix D).

Three- to five-hour observation periods are commonly recommended to document territorial occupancy of peregrine falcons and prairie falcons (USFWS 1984; Cade et al. 1996; Smith and Hutchins 2006). Steenhof et al. (1999) employed two-hour observation periods during point surveys to document territory occupancy of prairie and peregrine falcons in the Snake River region of Idaho. For a potential falcon territory to be classified as unoccupied at Pinnacles, we adopted a conservative standard of visiting potential nest sites at least three times per breeding season spaced 21–28 days apart to confirm territorial occupancy, courtship, and incubation of eggs within a breeding season (Fuller and Mosher 1981; Fraser et al. 1983; Steenhof 1998). Survey duration was ultimately dependent upon visibility, but at least three 4-hour surveys (12 hours total) were required to verify that “no birds” were present. Nests determined to be active were revisited to confirm rearing of nestlings and fledging of young. Nests in core areas were monitored more frequently and during weekend days when climbers were more likely to be present.

While other monitoring programs infer fledging success at 90% fledge age (Steenhof and Kochert 1982; Anderson and Squires 1997; Steenhof 1998), our protocol continues surveys until all young falcons are confirmed as fledged.

During the falcon breeding season status was asserted as follows:

Territories: Territorial behavior included perching, flying, territorial disputes and defense, stooping and scold calling, and roosting locations.

Courtship: Courtship behavior included copulation, food drops and swapping, and potential nest site inspections and preparation.

Incubation: Incubation status was determined by observing prairie falcons flying into a nest hole and not re-emerging for extended periods of time. During this time, egg counts were made whenever possible (e.g., when lighting conditions allowed and when incubating falcons temporarily left the nest during food drops and/or nest switches). Soft incubation – the onset of incubation – was determined by a small number of eggs laid and the female incubating for short durations (15–75 minutes of incubation and 20 minutes or more not incubating the eggs). Hard incubation was characterized by the adult falcons – primarily the females – incubating a full clutch of eggs for hours in duration.

Nestlings: Hatched young prairie and peregrine falcons were aged by physical features using an aging guide (Moritsch 1983). Hatch dates were determined by counting backwards from at least two (preferably three or more) independent aging estimates.

Fledging: Fledging was confirmed by seeing young perched and/or in flight away from the nest site. Fledging dates were estimated by the coordination and strength of flight, the size of perches, and the amount of vocalization during flight.

Monitoring Schedule

The prairie and peregrine falcon monitoring season started on 1 January and continued through the end of the nesting season, 13 July (Table 1).

Table 1. Timing of nesting behavior of prairie falcons at Pinnacles National Park.

Behavior	January	February	March	April	May	June	July
Territorial Falcons							
Courtship Behavior							
Nesting							
Fledging							

Weather was always an important factor. During temperature extremes, heavy fog, or rain, most birds of prey are generally inactive and therefore monitoring was not done during these periods.

Data Management

Data are entered into a Microsoft (MS) Access database designed by the network data manager for the San Francisco Bay Area Network Inventory and Monitoring Program. Original data sheets are archived with Pinnacles Resource Management. An annual (static) copy of the Access database is archived on the Golden Gate National Recreation Area computer network drive. Nest data are also submitted to the California Department of Fish and Wildlife (previously California Department of Fish and Game) California National Diversity Database.

Tabular data in the Results section of this report are derived from queries to the Breeding Raptors and Raptor Observations tables in the MS Access database.

Climbing Advisories

Climbing advisories went into effect by mid-January. Informational signs were established near territories occupied by prairie and peregrine falcons at least once during the preceding three years. Visitors were advised to avoid these areas but compliance was voluntary. Advisory areas with posted signs (Figure 2) included the Balconies, Hawkins, Scout Peak, and Little Pinnacles territories



Figure 2. Setting up advisory sign (NPS/GAVIN EMMONS, 2014).

Results

During the 2016 field season, Pinnacles staff spent 610 hours in the field surveying for prairie and peregrine falcons and volunteers contributed 100 hours of time. Results for prairie falcon monitoring in core and non-core areas are detailed below. Results for peregrine falcon monitoring and tabular data for combined prairie and peregrine falcon productivity are detailed in Appendix B.

Prairie Falcons

Six prairie falcon pairs attempted to nest this year and five successful nests produced 19 nestlings and fledglings, compared to 30-year averages of 9.9 nesting pairs, 7.8 successful nests, 28.9 nestlings, and 27.1 fledglings (Table 2).

Occupied Territories

Through the 2016 season, 11 territorial pairs of prairie falcons were confirmed over the course of the breeding season. This number is comparable to the average territorial occupancy of 11.9 pairs (1984–2015; Table 2). Of these, five pairs did not nest this year.

Core Areas: In 2016, there were seven territorial prairie falcon pairs within the core areas. The average number of territorial falcon pairs in the core areas over the previous 30 years was 7.2.

Non-Core Areas: In 2016 there were four territorial prairie falcon pairs within the non-core areas. The average number of territorial falcon pairs in the non-core areas over the last 30 years was 4.6.



Prairie falcon fledgling at South Balconies (NPS/GAVIN EMMONS, 2009).

Table 2. 1984–2016 Pinnacles prairie falcon nesting productivity – core and non-core areas combined.

Year	Territorial Pairs	Nesting Pairs	Successful Nests	# Nestlings	# Nestlings / Nest	# Fledglings	# Fledglings / Nest
1984	10	9	8	30	3.8	27	3.4
1987	6	4	4	13	3.3	10	2.5
1988	12	9	8	24	3.0	24	3.0
1989	12	12	9	24	2.7	21	2.3
1990	14	10	8	31	3.9	29	3.6
1991	14	11	10	34	3.4	34	3.4
1992	13	11	10	38	3.8	34	3.4
1993	13	12	10	39	3.9	35	3.5
1994	13	13	12	45	3.8	42	3.5
1995	13	11	8	24	3.0	24	3.0
1996	12	10	9	35	3.9	34	3.8
1997	12	8	6	26	4.3	26	4.3
1998	10	7	0	0	0	0	0
1999	10	8	6	25	4.2	25	4.2
2000	8	8	7	22	3.1	22	3.1
2001	10	10	7	24	3.4	24	3.4
2002	11	9	7	26	3.7	22	3.1
2003	12	9	8	33	4.1	32	4.0
2004	12	11	9	36	4.0	33	3.7
2005	13	10	9	29	3.2	24	2.7
2006	15	14	10	35	3.5	30	3.0
2007	14	12	9	35	3.9	33	3.7
2008	12	5	4	12	3.0	12	3.0
2009	12	11	10	41	4.1	37	3.7
2010	13	11	7	27	3.9	27	3.9
2011	13	12	8	33	4.1	33	4.1
2012	12	11	8	28	3.5	27	3.4
2013	12	11	10	47	4.7	43	4.3
2014	12	10	7	30	4.3	27	3.9
2015	11	8	5	21	4.2	21	4.2
2016	11	6	5	19	3.8	19	3.8
Averages (1984–2015)	11.9	9.9	7.8	28.9	3.6	27.1	3.4

Annual Productivity

Six of the 11 prairie falcon pairs nested. For the six nesting pairs, five successful nesting attempts fledged a total of 19 young (Tables 2, 3). One nest failed during the 2016 season during egg incubation.

Total nesting falcon pairs and successful nests within core and non-core areas this season were lower than the 30-year averages. Productivity of nestlings and fledglings (in total) were lower than the 30-year averages (Table 2, Figure 3).

Table 3. 2016 Pinnacles prairie falcon breeding summary.

Territory	Nest Used/ Last Year Used	# Eggs Laid	# Young Hatched	# Young Known/ Fledged
Crowley Towers*	CT-1/2014		5	5/5
Teapot Dome*	TD-3/2010	3	3	3/3
North Chalone	NC-7/NEW	4	4	4/4
Resurrection Wall*	RW-8/2015	3	3	3/3
South Balconies*	SGB-15/2015		4	4/4
Canyon North of Willow Spring Slide	CNWS-2/2006			0/0

*nests within the core area.




Figure 3. Core vs. non-core Pinnacles prairie falcon fledgling productivity, 1984–2016.

Core Areas: Of the seven territorial falcon pairs in the core areas in 2016, four nested successfully, producing 15 total fledglings and 3.8 fledglings per nest (Table 4). Productivity numbers are comparable to the 1984–2015 averages of 5.0 successful nests per season, 18.9 total fledglings, and 3.6 fledglings per nest.

Table 4. 1984–2016 Pinnacles prairie falcon nesting productivity – core areas only.

Year	Territorial Pairs	Nesting Pairs	Successful Nests	# Nestlings	# Nestlings / Nest	# Fledglings	# Fledglings / Nest
1984	7	6	6	22	3.7	20	3.3
1987	5	4	4	13	3.3	10	2.5
1988	8	6	5	14	2.8	14	2.8
1989	8	8	6	16	2.7	16	2.7
1990	9	7	6	23	3.8	23	3.8
1991	9	8	8	31	3.9	31	3.9
1992	9	7	7	29	4.1	27	3.9
1993	10	9	8	34	4.3	30	3.8
1994	10	10	10	38	3.8	37	3.7
1995	10	9	7	21	3.0	21	3.0
1996	9	8	7	28	4.0	24	3.4
1997	8	6	4	16	4.0	16	4.0
1998	7	5	0	0	0	0	0
1999	6	5	3	13	4.3	13	4.3
2000	5	5	4	15	3.8	15	3.8
2001	7	6	4	12	3.0	12	3.0
2002	5	5	4	12	3.0	12	3.0
2003	5	5	5	22	4.4	22	4.4
2004	7	7	5	21	4.2	18	3.6
2005	6	5	4	12	3.0	11	2.8
2006	7	6	5	17	3.4	15	3.0
2007	6	6	6	26	4.3	26	4.3
2008	7	3	2	4	2.0	4	2.0
2009	7	7	6	24	4.0	20	3.3
2010	8	6	3	11	3.7	11	3.7
2011	8	7	5	19	3.8	19	3.8
2012	7	6	3	14	4.7	14	4.7
2013	6	6	5	25	5.0	21	4.2
2014	6	4	4	16	4.0	16	4.0
2015	6	4	4	18	4.5	18	4.5
2016	7	4	4	15	3.8	15	3.8
Averages (1984– 2015)	7.2	6.2	5.0	18.9	3.6	17.9	3.4

Non-Core Areas: Of the four territorial falcon pairs in the non-core areas in 2016, one nested successfully, producing four total fledglings and 4.0 fledglings per nest (Table 5). These numbers are lower than the 1984–2015 average of 2.8 successful nests per season and 10.0 total fledglings, and higher than the 30-year average of 3.3 fledglings per nest.

Table 5. 1984–2016 Pinnacles prairie falcon nesting productivity – non-core areas only.

Year	Territorial Pairs	Nesting Pairs	Successful Nests	# Nestlings	# Nestlings / Nest	# Fledglings	# Fledglings / Nest
1984	3	3	2	8	4.0	7	3.5
1987	1	0	0	0	0	0	0
1988	4	3	3	10	3.3	10	3.3
1989	4	4	3	8	2.7	5	1.7
1990	5	3	2	8	4.0	6	3.0
1991	5	3	2	3	1.5	3	1.5
1992	4	4	3	9	3.0	7	2.3
1993	3	3	2	5	2.5	5	2.5
1994	3	3	2	7	3.5	5	2.5
1995	3	2	1	3	3.0	3	3.0
1996	3	2	2	7	3.5	10	5.0
1997	4	2	2	10	5.0	10	5.0
1998	3	2	0	0	0	0	0
1999	4	3	3	12	4.0	12	4.0
2000	3	3	3	7	2.3	7	2.3
2001	3	4	3	12	4.0	12	4.0
2002	6	4	3	14	4.7	10	3.3
2003	7	4	3	11	3.7	10	3.3
2004	5	4	4	15	3.8	15	3.8
2005	7	5	5	17	3.4	13	2.6
2006	8	8	5	18	3.6	15	3.0
2007	8	6	3	9	3.0	7	2.3
2008	5	2	2	8	4.0	8	4.0
2009	5	4	4	17	4.3	17	4.3
2010	5	5	4	16	4.0	16	4.0
2011	5	5	3	14	4.7	14	4.7
2012	5	5	5	14	2.8	13	2.6
2013	6	5	5	22	4.4	22	4.4
2014	6	6	3	14	4.7	11	3.7
2015	5	4	1	3	3.0	3	3.0
2016	4	2	1	4	4.0	4	4.0
Averages (1984–2015)	4.6	3.7	2.8	10.0	3.3	9.2	3.1

Phenology

The first prairie falcon pair was observed at Resurrection Wall on 1 January 2016 (Appendix A). Incubation was first observed at Resurrection Wall on 9 March. The first hatching occurred between 10 April and 15 April at Crowley Towers and the first fledging occurred between 22 May and 27 May at the same nest. The last fledging occurred at South Balconies between 10 June and 12 June.

Other Notes

Five of the six eyries chosen by prairie falcons were previously used. The North Chalone eyrie (located in a deep cavity near a cliff base) was previously undocumented. All prairie falcon eyries were within historically documented territories. One nesting attempt failed this year. Four territories occupied by prairie falcon pairs in the past five years – D. Soto Canyon, Pipsqueak Pinnacles, Narrows, and Marion Canyon – were vacant this year.

Discussion

Six prairie falcon pairs attempted to nest this year and five successful nests produced 19 nestlings and fledglings, compared to 30-year averages of 9.9 nesting pairs, 7.8 successful nests, 28.9 nestlings, and 27.1 fledglings (Table 2). In core areas in 2016, numbers for total successful nests, nestlings, and fledglings were comparable with 30-year averages (Table 4). In non-core areas, respective numbers were lower than average in 2016 (Table 5). One nest failure was documented during egg incubation at the single non-core area nest site in 2016. Given remoteness of the nest site, moderate temperatures during the period of nest failure, and common raven activity in the area, predation of eggs was suspected for the nest failure. However, given the lack of evidence or observations of predated eggs and lack of constant monitoring at the site (e.g., through remote video surveillance), other causes of nest failure cannot be entirely ruled out.

Prairie falcon productivity numbers for the 2016 season were lower than the 30-year averages of the Pinnacles raptor monitoring program.

Conclusions, Management Implications and Recommendations

Climbing management actions, outreach, and recommendations for further management and research are listed below. Refer to Appendix D for further information on public interest highlights for the 2016 season.

Prairie Falcons: Climbing Advisories

Climbing advisories were put in place in January in areas with historic climber usage to protect nesting raptors from disturbance. In March and April, advisories were updated and lifted in territories that were confirmed unoccupied by prairie and peregrine falcon pairs. Signs detailing climbing advisories were posted for Little Pinnacles, Balconies, Hawkins, Scout Peak, Crowley Towers, Egg, Tunnel, Teapot Dome, and Goat Rock / Resurrection Wall territories.

Due to the large size and climber popularity of Machete Ridge, a partial advisory was instituted at this territory. A partial advisory was also instituted at Balconies after a prairie falcon nest site was confirmed at South Balconies and a peregrine falcon nest site was confirmed at General Balconies. North Balconies was opened to climber use after the Balconies falcon pairs had shifted to focus territorial and nesting efforts at the General and South Balconies nest sites. The south end of Machete Ridge remained under advisory, and was then covered under a mandatory closure after July to protect a nesting California condor pair from climber and hiker disturbance.

All regular advisory signs were affixed to metal brackets and cement foundations to prevent theft. The General Balconies advisory sign was vandalized and replaced in 2014, but was not visibly tampered with in 2015 or 2016. No additional signs were disturbed or stolen.

In 2016, there were three incidents of off-trail hikers in advisory areas that were documented. There were also three incidents documented involving climbers in the Machete Ridge advisory area. No territorial defense behavior by prairie or peregrine falcons was observed by the hikers or falcon biologists during the off-trail hiker or climber incidents.

Human / Falcon Interactions and Nest Failures

Falcon adults in the North Chalone Peak, Crowley Towers, Teapot Dome, Resurrection Wall, and Balconies territories responded to the presence of on-trail hikers and raptor biologists with agitated behavior by circling and wailing above their respective territories.

Prairie falcon nest entries were conducted at the Crowley Towers, Resurrection Wall, North Chalone Peak, South Balconies, and Teapot Dome nests by the NPS raptor biologist, with additional assistance from the raptor monitoring intern (E. Cox), condor program crew members and biologists, and volunteers (see Public Interest Highlights below for further details). All nestlings at all 5 eyries fledged successfully.

A prairie falcon nest failure was documented at the Canyon North of Willow Spring territory. The nest failed during egg incubation. Given remoteness of the nest site, moderate temperatures during the period of nest failure, and common raven presence near (and subsequent nesting at) the falcon nest site, predation of eggs was suspected for the nest failure. Nest failure due to human disturbance, abandonment, or other causes cannot be entirely ruled out.

Education Opportunities

Throughout the year, the raptor biologist and park staff participated in public outreach opportunities to inform visitors about raptor conservation. Educational opportunities included participation in formal events (e.g., Climber Appreciation Days on 23–25 October, and scheduled PowerPoint presentations for visitors and staff at the East Side Search and Rescue Cache) and informal events (e.g., visitor contacts in high-use areas such as High Peaks, Balconies Cliff Trail, and the Bear Gulch Reservoir).

Management Recommendations

- Continue to establish climbing/hiking advisories in core areas (high visitor-use areas) each breeding season to protect cliff-nesting raptor species from human disturbance.
- Increase information opportunities for visitor use assistants and park rangers to educate park visitors about advisories. In past years (prior to 2004), park rangers and interpreters made more attempts to speak with climbers and hikers at trailheads and to regularly rove on trails to provide interpretation and enforcement of resources. Renewing this strategy of interfacing with visitors at trailheads and on trails would help to preserve compliance with climbing advisories as annual visitation increases at the park.
- Enforce advisories with law enforcement rangers. Although advisories are voluntary, disturbing wildlife is a citable offense that law enforcement rangers should continue to employ to discourage visitors from willfully threatening nesting efforts of breeding raptors at Pinnacles.

- Continue use of staff and visitors to observe raptor activity in the field. This can be achieved through regular communication with NPS staff and visitors, continued use of monthly monitoring updates on raptor status at the park, and reminders about filling out wildlife observation cards.

Literature Cited

- Anderson, S. H., and J. R. Squires. 1997. The prairie falcon. University of Texas Press. Austin, Texas.
- Becker, D. M., and I. J. Ball. 1981. Impacts of surface mining on prairie falcons: Recommendations for monitoring and mitigation. Montana Cooperative Wildlife Research Unit Unpublished Report, Montana State University, Missoula.
- Bednarz, J. C. 1984. Effect of mining and blasting on breeding prairie falcon (*Falco mexicanus*) occupancy in the Caballo Mountains, New Mexico. Raptor Research 18:16–19.
- Bloom, P. H. 1994. The biology and current status of the long-eared owl in coastal southern California. Bulletin California Academy of Science 93:1–12.
- Bond, R. M. 1946. The peregrine populations of western North America. Condor 48:101–116.
- Boyce, D. A., Jr. 1982. Prairie falcon fledgling productivity in the Mojave Desert, California. Thesis. Humboldt State University, Arcata, California.
- Cade, T. J., J. H. Enderson, and J. Linthicum. 1996. Guide to management of peregrine falcons at the eyrie. The Peregrine Fund, Boise, Idaho.
- Delannoy, C. A., and A. Cruz. 1988. Breeding biology of the Puerto Rican sharp-shinned hawk (*Accipiter striatus venator*). Auk 105:649–662.
- Emmons, G., J. Pettersen, M. Koenen, and D. Press. 2011. Prairie falcon monitoring protocol for Pinnacles National Monument: narrative – version 2.3. Natural Resource Report NPS/SFAN/NRR—2011/466. National Park Service, Fort Collins, Colorado.
- Fraser, J. D., L. D. Frenzel, J. E. Mathisen, F. Martin, and M. E. Shough. 1983. Scheduling bald eagle reproduction surveys. Wildlife Society Bulletin 11:13–16.
- Fuller, M. R., and J. A. Mosher. 1981. Methods of detecting and counting raptors: a review. Studies in Avian Biology 6:235–246.
- Fyfe, R. W., and R. R. Olendorff. 1976. Minimizing the dangers of nesting studies to raptors and other sensitive species. Canadian Wildlife Service Occasional Papers No. 23:1–17.
- Harmata, A. R., J. E. Durr, and H. Geduldig. 1978. Home range, activity patterns and habitat use of Prairie Falcons nesting in the Mojave Desert. Colorado Wildlife Services Unpublished Report, Fort Collins, Colorado for U.S. Department of the Interior, Bureau of Land Management, Riverside, California (Contract No. YA-512-CT8-4389).
- Hickey, J. J. 1942. Eastern population of the duck hawk. Auk 59: 176–204.

- Hickey, J. J. 1969. Peregrine falcon populations: their biology and decline. University of Wisconsin Press, Madison, Wisconsin.
- Holthuijzen, A. M. A., W. G. Eastland, A. R. Ansell, M. N. Kochert, R. D. Williams, and L. S. Young. 1990. Effects of blasting on behavior and productivity of nesting prairie falcons. *Wildlife Society Bulletin* 18:270–281.
- Kochert, M. N., K. Steenhof, L. B. Carpenter, and J. M. Marzluff. 1999. Effects of fire on golden eagle territory occupancy and reproductive success. *Journal of Wildlife Management* 63:773–780.
- Marks, J. S. 1986. Nest-site characteristics and reproductive success of long-eared owls in southwestern Idaho. *Wilson Bulletin* 98:547–560.
- Marti, C. D., and J. S. Marks. 1989. Medium-sized owls. *Proceedings of the Western Raptor Management Symposium and Workshop*. National Wildlife Federation Science and Technology Series No. 12, Washington, D.C.:124–133.
- Moritsch, M. Q. 1983. Photographic guide for aging nesting prairie falcons. United States Department of the Interior, Bureau of Land Management, Snake River Birds of Prey Project. Boise District, Idaho.
- Newton, I. 1979. Population ecology of raptors. Buteo Books. Shipman, Virginia.
- Newton, I. 1990. Birds of prey. Facts on File, Inc. New York, New York.
- Ogden, V. T., and M. G. Hornocker. 1977. Nesting density and success of prairie falcons in southwestern Idaho. *Journal of Wildlife Management* 41:1–11.
- Olsen, P., and J. Olsen. 1978. Alleviating the impact of human disturbance on the breeding peregrine falcon. *Ornithologists Corella* 2(1):1–7 and 4(3):54–57.
- Platt, S. W. 1974. Breeding status and distribution of the prairie falcon in northern New Mexico. Thesis. Oklahoma State University, Stillwater, Oklahoma.
- Porter, R. D., M. A. Jenkins, and A. L. Ganski. 1987. Working bibliography of the peregrine falcon. National Wildlife Federation Science and Technology Series No. 9, Washington, D.C.
- Scott, T. A. 1985. Human impacts on the golden eagle population of San Diego County. Thesis. San Diego State University, San Diego, California.
- Sitter, G. 1983. Feeding activity and behavior of prairie falcons in the Snake River Birds of Prey Natural Area in southwestern Idaho. Thesis. University of Idaho, Moscow, Idaho.
- Smith, J. P., and A. Hutchins. 2006. Northeast Nevada Nest Survey 2005. Hawkwatch International, Inc., Salt Lake City, Utah. Available from

<http://www.hawkwatch.org/publications/Technical%20Reports/NNNS%20Report%202005.pdf>
(accessed 20 September 2008).

Steenhof, K. 1998. Prairie falcon (*Falco mexicanus*). The Birds of North America, No. 346. Cornell Laboratory of Ornithology, Ithaca, New York.

Steenhof, K., and M. N. Kochert. 1982. An evaluation of methods used to estimate raptor nesting success. *Journal of Wildlife Management* 46:885–893.

Steenhof, K., M. N. Kochert, L. B. Carpenter, and R. N. Lehman. 1999. Long-term prairie falcon population changes in relation to prey abundance, weather, land uses, and habitat conditions. *Condor* 101:28–41.

Steenhof, K., M. N. Kochert, and T. L. McDonald. 1997. Interactive effects of prey and weather on golden eagle reproduction. *Journal of Animal Ecology* 66:350–362.

Steidl, R. J., K. D. Kozie, G. J. Dodge, T. Pehovski, and E. R. Hogan. 1993. Effects of human activity on breeding behavior of golden eagles in Wrangell-St. Elias National Park and Preserve; a preliminary assessment. WRST Research and Resource Management Report No. 93-3. National Park Service, Wrangell-St. Elias National Park and Preserve, Copper Center, Alaska.

Suter, G. W., and J. L. Jones. 1981. Criteria for golden eagle, ferruginous hawk, and prairie falcon nest site protection. *Raptor Research* 15:12–18.

United States Department of the Interior (USDI). 1979. Snake River Birds of Prey special research report to the Secretary of the Interior. Boise District Bureau of Land Management, Boise, Idaho.

United States Fish and Wildlife Service (USFWS). 1984. American peregrine falcon Rocky Mountain/Southwest population recovery plan. Rocky Mountain/Southwest Peregrine Falcon Recovery Team, U.S. Fish and Wildlife Service, Denver, Colorado.

Watson, J. 1997. The golden eagle. London, United Kingdom.

Western Regional Climate Center (WRCC). 2013. Climate data for Pinnacles National Monument. Available from <http://www.wrcc.dri.edu/> (accessed August 2013).

White, C. M., N. J. Clum, T. J. Cade, and W. G. Hunt. 2002. Peregrine falcon (*Falco peregrinus*). A. Poole, editor. The birds of North America online. Cornell Laboratory of Ornithology, Ithaca, New York. Available at http://bna.birds.cornell.edu/BNA/account/Peregrine_Falcon (accessed 8 September 2008).

Appendix A. 2016 nest phenology and success for prairie and peregrine falcons.

Nest Species	Territory Occupied	Nest Code	Arrival Date	Begin Incub	Hatch Date	Fledge Date	Abandon Date	Failed Date	# of Eggs	# of Nestlings	Known Fledglings	Possible Fledglings	Occup. Status
PRFA	Ball Pinnacle	*TD-3	<1/20										Occupied
PRFA	Cyn N of Willow Spgs	CNWS-2	<1/15	<4/4				<5/11			0	0	Failed
PRFA	Central High Peaks												Not Occ.
PRFA	Citadel		1/20-26				>2/12						Abandon
PRFA	Crowley Drainage												Not Occ.
PRFA	Crowley Towers	CT-1	<1/15	3/20-31	4/10-15	5/22-27				5	5	5	5
PRFA	D. Soto Canyon												Not Occ.
PRFA	Deserted Valley												Not Occ.
PRFA	Discovery Wall												Not Occ.
PRFA	Drywall		<3/24										Occupied
PRFA	Egg	*TD-3	<1/20										Occupied
PRFA	Frog Canyon												Not Occ.
PRFA	Frog / Hand		<3/9				>3/9						Abandon
PRFA	General Balconies												Not Occ.
PRFA	Goat Rock	*RW-8	<1/28										Occupied
PRFA	Guard Rock												Not Occ.
PRFA	Hanging Valley												Not Occ.
PRFA	Hawkins												Not Occ.
PRFA	High Pks W of CPA												Not Occ.
PRFA	Little Pinnacles		<1/12				>1/12						Abandon
PRFA	Machete												Not Occ.
PRFA	Marion Canyon												Not Occ.
PRFA	Mating Rocks												Not Occ.
PRFA	Narrows												Not Occ.
PRFA	NE Sec 15												Not Occ.

Note: for the "Occup. Status" column, # refers to possible fledglings, "Occupied" = territorial occupation, "Not Occ." = no occupation, "Failed" = failed nest, "Abandon" = territory abandoned after confirmed occupancy, "Unknown" = breeding confirmed (see nest code) or likely, but nest status unknown. For the "Nest Code" column, * refers to territorial links for raptor pairs occupying more than 1 territory.

Appendix A (continued). 2016 nest phenology and success for prairie and peregrine falcons.

Nest Species	Territory Occupied	Nest Code	Arrival Date	Begin Incub	Hatch Date	Fledge Date	Abandon Date	Failed Date	# of Eggs	# of Nestlings	Known Fledglings	Possible Fledglings	Occup. Status
PRFA	Neglected Valley												Not Occ.
PRFA	North Balconies												Not Occ.
PRFA	North Chalone	NC-7	<1/23	<3/23	4/21-23	5/29-6/3				4	4	4	4
PRFA	North Wilderness Rock												Not Occ.
PRFA	Pig Canyon												Not Occ.
PRFA	Pipsqueak Pinnacles												Not Occ.
PRFA	Prescribed Burn Cliffs												Not Occ.
PRFA	Resurrection Wall	RW-8	<1/1	<3/9	4/15-19	5/26-30			3	3	3	3	3
PRFA	Scout Peak	*RW-8	<1/1										Occupied
PRFA	South Balconies	SGB-15	<1/20	<4/25	4/29-5/1	6/10-12				4	4	4	4
PRFA	South Chalone		<3/23										Occupied
PRFA	S. Wilderness Rock												Not Occ.
PRFA	Teapot Dome	TD-3	<1/20	<3/30	4/21-24	6/2-5			3	3	3	3	3
PRFA	Tugboat												Not Occ.
PRFA	Tunnel	*TD-3	<1/20										Occupied
PRFA	Upper Bear Gulch												Not Occ.
PRFA	Upper Condor Gulch												Not Occ.
PRFA	Western Front												Not Occ.
PRFA	Willow Spring Slide	*CNWS-2	<1/15										Occupied
PEFA	Hawkins Peak	HP-1	<1/1	3/30-4/13				5/5-12	3	0	0	0	Failed
PEFA	Upper Condor Gulch	*HP-1	<1/1										Occupied
PEFA	Central High Peaks	*HP-1	<1/1										Occupied
PEFA	General Balconies	SGB-6	<1/20	3/21-28	5/1-3	6/12-16			3	3	3	3	3
PEFA	Machete	*SGB-6	<1/20										Occupied
PEFA	North Balconies	*SGB-6	<1/20										Occupied
PEFA	Drywall												Not Occ.

Note: for the "Occup. Status" column, # refers to possible fledglings, "Occupied" = territorial occupation, "Not Occ." = no occupation, "Failed" = failed nest, "Abandon" = territory abandoned after confirmed occupancy, "Unknown" = breeding confirmed (see nest code) or likely, but nest status unknown. For the "Nest Code" column, * refers to territorial links for raptor pairs occupying more than 1 territory.

Appendix B. 2016 results and discussion for peregrine falcon occupancy and productivity.

Results: Peregrine Falcons

Two peregrine falcon pairs occupied territories at Pinnacles in 2016 and nested at Hawkins Peak and General Balconies (Table APP B.1). This marked the twelfth consecutive year that a peregrine falcon pair has nested at Pinnacles. Prior to 2004, peregrine falcon occupancy or nesting had not been confirmed at Pinnacles for 48 years. The Hawkins nest effort failed during egg incubation. The General Balconies pair fledged 3 young (Figure APP B.1), compared to 12-year averages (since 2004) of 1.1 nesting pairs, 1.0 successful nests, 2.8 nestlings, and 2.3 fledglings (Table APP B.2).

Occupied Territories

Core Areas: In 2016 there were two territorial peregrine falcon pairs within the core areas. This number is higher than the average number of territorial peregrine falcon pairs (1.3) in the core areas over the last ten years.

Non-Core Areas: In 2016 there were no territorial falcon pairs within non-core areas.

Phenology and Productivity

The peregrine falcon pairs at Hawkins Peak and General Balconies were first observed this year on 1 and 20 January 2016, respectively (Appendix A), but were also documented occupying and defending the respective territories throughout the fall and winter of 2015. Incubation was first observed by the General Balconies peregrine falcon pair on 21–28 March. Hatching of young occurred at the General Balconies nest on 1–3 May. Fledging took place at the General Balconies nest on 16–18 June.



Figure APP B.1. Peregrine falcon fledgling at Hawkins Peak (NPS/GAVIN EMMONS, 2012).

Table APP B.1. 2016 Pinnacles peregrine falcon breeding summary.

Territory	Nest Used/ Last Year Used	# Eggs Laid	# Young Hatched	# Young Known/ Fledged
Hawkins Peak *	HP-1/ 2013		0	0/0
General Balconies *	SGB-6/ 2015	3	3	3/3

*nests within the core area.

Table APP B.2. 1984–2016 Pinnacles peregrine falcon nesting productivity—core areas only.

Year	Territorial Pairs	Nesting Pairs	Successful Nests	# Nestlings	# Nestlings / Nest	# Fledglings	# Fledglings / Nest
1984	0	0	0	0	0	0	0
1987	0	0	0	0	0	0	0
1988	0	0	0	0	0	0	0
1989	0	0	0	0	0	0	0
1990	0	0	0	0	0	0	0
1991	0	0	0	0	0	0	0
1992	0	0	0	0	0	0	0
1993	0	0	0	0	0	0	0
1994	0	0	0	0	0	0	0
1995	0	0	0	0	0	0	0
1996	0	0	0	0	0	0	0
1997	0	0	0	0	0	0	0
1998	0	0	0	0	0	0	0
1999	0	0	0	0	0	0	0
2000	0	0	0	0	0	0	0
2001	0	0	0	0	0	0	0
2002	0	0	0	0	0	0	0
2003	0	0	0	0	0	0	0
2004	1	0	0	0	0	0	0
2005	1	1	1	3	3	3	3
2006	1	1	1	3	3	3	3
2007	1	1	1	3	3	3	3
2008	1	1	1	3	3	3	3
2009	1	1	1	3	3	3	3
2010	1	1	1	1	1	1	1
2011	1	1	0	0	0	0	0
2012	1	1	1	4	4	4	4
2013	2	1	1	4	4	3	3
2014	2	2	2	3	1.5	3	1.5
2015	2	2	2	6	3	6	3
2016	2	2	1	3	3	3	3
Averages (2004–2015)	1.3	1.1	1.0	2.8	2.4	2.7	2.3

Discussion: Combined Prairie and Peregrine Falcon Occupancy and Productivity

Given the small sample size and brief period of years that peregrine falcon pairs have occupied and produced young at breeding territories at Pinnacles, we cannot currently assign any statistical relevance or trend analyses specifically to peregrine falcon breeding results at the park. However, we have combined prairie and peregrine falcon results over the 30 years of monitoring efforts (Table APP B.3 and Figure APP B.2) to represent overall large falcon occupancy and productivity numbers at Pinnacles.

Combined large falcon occupancy and productivity in core and non-core areas this year were generally lower than the 30-year running average rates. Eight falcon pairs attempted to nest this year and 6 successful nests produced 22 nestlings and fledglings, compared to 30-year averages of 10.3 nesting pairs, 8.2 successful nests, 30.0 nestlings, and 28.1 fledglings (Table APP B.3).

In comparing prairie falcon results (Table 2 and Figure 3) with combined prairie and peregrine falcon results (Table APP B.3 and Figure APP B.2), peregrine falcon breeding efforts appear to be supplementing overall large falcon productivity to yield more stable numbers over the 30 years of falcon monitoring at Pinnacles. A prairie falcon pair consistently nested at Hawkins Peak before a peregrine pair began occupying the territory 10 years ago; since 2004 no prairie falcons have nested at Hawkins Peak. This may suggest that – in certain instances – prairie and peregrine falcons compete for territorial occupancy of limited cliff-nest habitat at Pinnacles, with total large falcon productivity unaffected but lower total productivity for prairie falcons in the future, particularly if additional peregrine falcon pairs re-occupy more historical territories. However, in 2014 to 2016 prairie and peregrine falcon pairs in relatively close proximity—at South and General Balconies—both successfully nested and fledged young. The prairie and peregrine falcon pairs at Balconies defended the cliff areas and limited confrontations between the falcon pairs were observed. At present, limited peregrine nesting efforts and data are inconclusive in regard to competition for cliff-nest habitat between the two species.

Table APP B.3. 1984–2016 Pinnacles prairie and peregrine falcon nesting productivity – core and non-core areas combined.

Year	Territorial Pairs	Nesting Pairs	Successful Nests	# Nestlings	# Nestlings / Nest	# Fledglings	# Fledglings / Nest
1984	10	9	8	30	3.8	27	3.4
1987	6	4	4	13	3.3	10	2.5
1988	12	9	8	24	3.0	24	3.0
1989	12	12	9	24	2.7	21	2.3
1990	14	10	8	31	3.9	29	3.6
1991	14	11	10	34	3.4	34	3.4
1992	13	11	10	38	3.8	34	3.4
1993	13	12	10	39	3.9	35	3.5
1994	13	13	12	45	3.8	42	3.5
1995	13	11	8	24	3.0	24	3.0
1996	12	10	9	35	3.9	34	3.8
1997	12	8	6	26	4.3	26	4.3
1998	10	7	0	0	0	0	0
1999	10	8	6	25	4.2	25	4.2
2000	8	8	7	22	3.1	22	3.1
2001	10	10	7	24	3.4	24	3.4
2002	11	9	7	26	3.7	22	3.1
2003	12	9	8	33	4.1	32	4.0
2004	13	11	9	36	4.0	33	3.7
2005	14	11	10	32	3.2	27	2.7
2006	16	15	11	38	3.5	33	3.0
2007	15	13	10	38	3.8	36	3.6
2008	13	6	5	15	3.0	15	3.0
2009	13	12	11	44	4.0	40	3.6
2010	14	12	8	28	3.5	28	3.5
2011	14	13	8	33	4.1	33	4.1
2012	13	12	9	32	3.6	31	3.4
2013	14	12	11	51	4.6	46	4.2
2014	14	12	9	33	3.7	30	3.3
2015	14	10	7	27	3.9	27	3.9
2016	13	8	6	22	3.7	22	3.7
Averages (1984–2015)	12.4	10.3	8.2	30.0	3.5	28.1	3.3




Figure APP B.2. Combined core and non-core Pinnacles PRFA and PEFA fledgling productivity, 1984–2016.

Appendix C. Documentation of changes in data collection methods.

No changes were made to data collection methods for the Pinnacles raptor monitoring program through the course of the 2016 season.

Following the end of the 2016 field season, the front end of the Raptor Monitoring Access database was updated with an improved Data Entry Form interface and removed several redundant fields. The back end of the database was updated with additional Data Summary Tools including improved quality assurance (QA) queries for duplicate and missing records, and for checking errors in survey locations and falcon occupancy determinations. The Breeding Raptors table was also updated for improved and more consistent terminology in occupancy status fields.

Following the end of the 2015 field season, legacy data were revised to account for negative data (i.e., no birds observed) for all prairie and peregrine falcon observations for the years of 2003–2015. Legacy data were also revised to account for missing data in records including observer location, weather, and falcon territories observed.

Following the end of the 2014 field season, the Raptor Monitoring Access database was updated from .mdb to .accdb format. Terminology and data entry fields in the Data Entry Form were updated to more clearly reflect raptor observations pertaining to the I&M Protocol versus observations of other raptor species in the park, to distinguish more efficiently between occupancy and fecundity results, and to provide easier access to territory tables. Data summary tools were also added as queries to allow for more comprehensive review and revision of breeding summary and raptor observation information in spreadsheet format.

Appendix D. Public interest highlights.

The 2016 breeding season was the 31st year of raptor monitoring at Pinnacles. Field observations began 1 January 2016 and ended 1 July 2016, with a total of over 150 possible and active nest sites monitored during 710 observation hours. Climbing advisories were put into effect in January to reduce nest disturbance by visitors, updated to reflect unoccupied territories in March, and lifted in late July at the end of the raptor breeding season.

- During the 2016 season, the NPS raptor biologist independently conducted prairie falcon nest entries at five Pinnacles prairie falcon eyries after gaining Bird Banding Lab sub-permittee status (in 2013) under the master banding permit of raptor researcher Dr. Doug Bell from East Bay Regional Park District. All falcon nestlings at the Crowley Towers, Resurrection Wall, North Chalone Peak, South Balconies, and Teapot Dome nests were briefly handled, banded, and blood samples were obtained. All nestlings from the five eyries fledged successfully. Blood samples taken from all nestlings were screened for rodenticide exposure; test results are still forthcoming. (Three nests were sampled in 2014, and two more nests were sampled in 2015. During the 2014 season, all four falcon nestlings at the South Balconies nest tested positive for trace amounts of Diphacinone, representing the first confirmation of rodenticide exposure in the Pinnacles prairie falcon population.) Additionally, banding and blood sample information collected in 2009–2016 will be used to ascertain genetic insularity and pair fidelity in the Pinnacles prairie falcon population.
- In November 2015, the raptor biologist co-presented (with condor field crew supervisor Jennie Jones) at the 2015 Raptor Research Foundation conference in Sacramento. The 20-minute presentation focused on preliminary results of rodenticide exposure in prairie falcons and California condors (*Gymnogyps californianus*) and the challenges of managing breeding falcon and condor populations at Pinnacles National Park in the context of toxins that both species are exposed to in the surrounding working landscapes.
- During the 2016 season, one territorial pair of peregrine falcons successfully fledged a total of 3 young, representing the twelfth consecutive year that a pair of peregrine falcons has successfully nested at the park during the history of the raptor monitoring program. Prior to 2004, the last previously confirmed peregrine falcon territorial occupancy or nest effort at Pinnacles was documented at the Drywall territory in 1957.
- An ArcMap project was updated to visually display GPS and GIS information relating to the raptor monitoring program, including historical nest sites, monitoring watch spots, nest distribution by geologic and habitat layers, and locations of advisory signs posted at Pinnacles.
- PowerPoint presentations focused on raptor monitoring and falcon nesting were given to incoming Pinnacles interpretive staff and volunteers.
- The Third Annual Pinnacles Climber Appreciation Days (PCAD) event was organized by John Donham (the acting Pinnacles trails foreman), M. A. K. Spero (law enforcement ranger), Larry

Arthur (owner of Mountain Tools and a local climber), and the Pinnacles raptor biologist. During the three-day event, local climbers volunteered to restore and maintain climber access trails at popular climbing formations in Bear Gulch. The raptor biologist also used informal interpretive opportunities to discuss resource management, falcon cliff-nesting, and climbing advisories with the 70+ volunteers.

The Department of the Interior protects and manages the nation's natural resources and cultural heritage; provides scientific and other information about those resources; and honors its special responsibilities to American Indians, Alaska Natives, and affiliated Island Communities.

NPS 114/136151, January 2017

National Park Service
U.S. Department of the Interior



Natural Resource Stewardship and Science
1201 Oakridge Drive, Suite 150
Fort Collins, CO 80525

www.nature.nps.gov

EXPERIENCE YOUR AMERICA™